laticollis Wagner under the name of Aethus pilosulus Klug (vide Wagner 1954, pp. 2-3).

ACKNOWLEDGEMENTS

I take this opportunity to express my thanks to Dr. P. Freeman, Keeper, Department of Entomology, British Museum (Natural History), London, for facilities provided. Thanks are also due to Dr. G. S. Sandhu who supplied the material.

Commonwealth Institute of Entomology, M. S. K. GHAURI 56, Queen's Gate, London, S.W.7, February 17, 1972.

REFERENCES

AHMAD, T. & GHAURI, A. S. K. (1953): Studies on Acheta domestica (Linn.) I. Biology and Habits. Agric. Pakistan 4:235-286.

Dallas, W. S. (1851): List of the specimens of hemipterous insects in the collection of the British Museum. Pt. 1., 368 p., 11 plus.

STICHEL, W. (1961): Illustriete Bes-

timmungstabellen der Wanzen. II Europa (Hemiptera-Heteroptera Europae) (Cydnidae) vol. 4, 22 heft: 688.

WAGNER, E. (1954): Neue Heteropteren von den Kanarischen Inseln. Commentat. biol. (1) 14:1-28.

WESTWOOD, J. O. (1837): Catalogue of Hemiptera in the collection of the Rev. F. W. Hope. London.

24. DATURA SUAVEOLENS H.B. EX WILLD. (SOLANACEAE)—FROM DISTRICT CHAMOLI IN WESTERN HIMALAYAS

Datura suaveolens H.B. ex. Willd. Enum. Hort. Berol. 227.

An indigenous species of Mexico introduced in India as an ornamental plant. Raizada (1931) recorded it, growing in shady places in perfectly naturalized conditions, from Dehradun of upper Gangetic plain. In 1936 Raizada further gave its account.

During a floristic field study of Karanprayag block in Chamoli district I collected this species, from village Jakh, Tallachandpur, near Manda Khali on 24.vi.72. *Nautiyal* 5502, 145 m; wild in dry exposed habitat.

The taxon is distinguished by its shrubby habit and large, pendulous, white, sweet scented, flowers.

DEPARTMENT OF BOTANY, MEERUT UNIVERSITY, MEERUT, June 18, 1973.

K. N. NAUTIYAL